

Sleep and adaptation disorders in military personnel with blast injury

Kateryna Sarazhyna , O Ruskykh, I Serbin, Y Solodovnikova, A Son

Sleep and adaptation disorders are common in traumatic brain injury (TBI) resulting from blast injuries (BIs) and can significantly complicate the recovery, leading to delayed neurocognitive, psychiatric and behavioural disorders.^{1,2} Nevertheless, auditory disorders and tinnitus after repeated TBIs significantly worsen and affect daily activity through impaired sleep quality, irritability, anxiety and difficulties with concentration.³⁻⁵ We conducted this study to analyse the characteristics of sleep and adaptation disorders in military personnel with BIs.

Data were collected from October 2023 to April 2024 among 35 military personnel with BIs aged between 22 and 56 years (mean=31, SD=33.9), undergoing treatment at the University Clinic, with the exclusion of patients who had sleep disorders before the injury and included only military personnel who had sustained a mild TBI and were in the acute phase of their recovery. Sleep quality was assessed using the modified Pittsburgh Sleep Quality Index (PSQI). A sleep quality score of 0–7 points was considered satisfactory, scoring above 7 points was categorised as having poor sleep quality. Adaptation disorders were identified using the Hospital Anxiety and Depression Scale (HADS). On the HADS, scores above 11 were considered significant and indicative of clinical manifestations of adaptation disorder. Participants were divided into three groups: group 1—BI with TBI, barotrauma and tinnitus (n=5, 14.3%); group 2—BI with TBI without tinnitus (n=14, 40%); group 3—control group, BI without TBI (n=16, 45.7%).

Table 1 presents summary data from a study involving three separate groups, detailing measures of anxiety, depression and sleep quality.

The Shapiro-Wilk test was applied to assess the normality of data distribution to three variables (HADS anxiety, HADS depression, PSQI) across the groups, revealing: group 1: HADS anxiety

Table 1 Descriptive analysis

Groups	Mean HADS anxiety	SD HADS anxiety	Mean HADS depression	SD HADS depression	Mean PSQI	SD PSQI	Mean age	Percentage
1	11.60	5.37	15.20	2.39	14.80	1.30	35.40	14.29
2	8.00	3.55	9.07	3.85	11.36	2.71	36.07	40.00
3	6.50	2.92	6.81	3.47	6.69	2.15	31.50	45.71

HADS, Hospital Anxiety and Depression Scale; PSQI, Pittsburgh Sleep Quality Index.

Table 2 Post hoc tests

Comparison	Mean difference HADS anxiety	P values (Tukey)	Mean difference HADS depression	P values (Tukey)	Mean difference PSQI	P values (Tukey)
Group 1 vs group 2	3.60	0.145	6.13	0.006	3.44	0.020
Group 1 vs group 3	5.10	0.023	8.39	<001	8.11	<001
Group 2 vs group 3	1.50	0.492	2.26	0.201	4.67	<001

HADS, Hospital Anxiety and Depression Scale; PSQI, Pittsburgh Sleep Quality Index.

p=0.502; HADS depression p=0.294; PSQI p=0.421. Group 2: HADS anxiety p=0.192; HADS depression p=0.909; PSQI p=0.091. Group 3: HADS anxiety p=0.114; HADS depression p=0.628; PSQI p=0.896. We used analysis of variance to compare the quality among the groups that fit the normal distribution profile. HADS Anxiety: F-statistic=3.92, p=0.030. HADS Depression: F-statistic=10.85, p=0.00025. PSQI Sleep Quality: F-statistic=29.27, p=5.92e-08. To determine the differences between specific groups, we used post hoc tests, specifically the Tukey test (table 2).

The results confirm that tinnitus correlates with a significant increase in anxiety and depression with significant differences in sleep quality between the groups (F-statistic=29.27, p<0.001). A potential negative impact of tinnitus on the psychological state of military personnel who have sustained TBI has been identified. Tinnitus can be considered an independent factor in worsening sleep quality and adaptation. The main limitation of our study is the relatively small sample size (n=35) and the inclusion of only men in the study group. This suggests that that should be considered when assessing the condition of patients with various forms of BI.

X Kateryna Sarazhyna @katjasara

Contributors KS is the main author and guarantor responsible for writing the manuscript. OR performed the calculations and assisted in writing the manuscript. IS conducted data collection and contributed to analysing and interpreting data for the work. YS agreed to be accountable for all aspects of the work, ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. AS provided final approval for the published version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Local Ethics Committee (Odessa National Medical University) N 0842/27 dated 14 March 2024. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; internally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Neurology and Neurosurgery, Odessa National Medical University, Odessa, Ukraine

Correspondence to Kateryna Sarazhyna; katarazina@gmail.com



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/military-2024-002793>).



To cite Sarazhyna K, Ruskykh O, Serbin I, *et al*. *BMJ Mil Health* Epub ahead of print: [please include Day Month Year]. doi:10.1136/military-2024-002793

Received 26 May 2024

Accepted 20 August 2024

BMJ Mil Health 2024;**0**:1–2.

doi:10.1136/military-2024-002793

ORCID iD

Kateryna Sarazhyna <http://orcid.org/0000-0003-3332-1743>

REFERENCES

- 1 Sandsmark DK, Elliott JE, Lim MM. Sleep-Wake Disturbances After Traumatic Brain Injury: synthesis of Human and Animal Studies. *Sleep* 2017;40:zsx044.
- 2 Osborn AJ, Mathias JL, Fairweather-Schmidt AK. Prevalence of anxiety following adult traumatic brain injury: a meta-analysis comparing measures, samples and postinjury intervals. *Neuropsychology* 2016;30:247–61.
- 3 Shao N, Skotak M, Pendyala N, *et al*. Temporal Changes in Functional and Structural Neuronal Activities in Auditory System in Non-Severe Blast-Induced Tinnitus. *Medicina (Kaunas)* 2023;59:1683.
- 4 Singh AK, Ditkofsky NG, York JD, *et al*. Blast Injuries: from Improvised Explosive Device Blasts to the Boston Marathon Bombing. *Radiographics* 2016;36:295–307.
- 5 Swan AA, Nelson JT, Swiger B, *et al*. Prevalence of hearing loss and tinnitus in Iraq and Afghanistan Veterans: a Chronic Effects of Neurotrauma Consortium study. *Hear Res* 2017;349:4–12.